

BITSAT

Birla Institute of Technology & Science Aptitude Test

Mock Test -1

PHYSICS

1. In two system of unit relations among velocity acceleration and force are respectively $v_2 = \frac{\alpha^2}{\beta} v_1$, $a_2 = \alpha\beta a_1$, and $F_2 = \frac{F_1}{\alpha\beta}$. If α and β are constant then relations among mass, length and time in two system are -

(a) $M_2 = \frac{\alpha}{\beta} M_1$, $L_2 = \frac{\alpha^2}{\beta^2} L_1$, $T_2 = \frac{\alpha^3 T_1}{\beta}$

(b) $M_2 = \frac{1}{\alpha^2 \beta^2} M_1$, $L_2 = \frac{\alpha^3}{\beta^3} L_1$, $T_2 = T_1 \frac{\alpha}{\beta^2}$

(c) $M_2 = \frac{\alpha^3}{\beta^3} M_1$, $L_2 = \frac{\alpha^2}{\beta^2} L_1$, $T_2 = \frac{\alpha}{\beta} T_1$

(d) $M_2 = \frac{1}{\alpha^2 \beta^2} M_1$, $L_2 = \frac{\alpha}{\beta^2} L_1$, $T_2 = \frac{\alpha^3}{\beta^3} T_1$

2. What is the maximum range that a ball thrown with a speed of 40 ms^{-1} can cover without hitting the 25 m high ceiling of a long hall ?

- (a) 150.5 m (b) 100.25 m
 (c) 110.3 m (d) 200.5 m

3. If $\vec{P} + \vec{Q} + \vec{R} = 0$ and out of these, two vectors are equal in magnitude and the third vector has magnitude $\sqrt{2}$ times that of any of these two vectors, then angles among the three vectors are -

- (a) $45^\circ, 75^\circ, 75^\circ$ (b) $45^\circ, 90^\circ, 135^\circ$
 (c) $90^\circ, 135^\circ, 180^\circ$ (d) $90^\circ, 135^\circ, 135^\circ$


4. A body takes n times, the time to slide down a rough inclined plane as it takes to slide down the same inclined plane when it is perfectly frictionless. The coefficient of kinetic friction between body and the plane for an angle of inclination of 45° is given by μ -

- (a) $1 - \frac{1}{n}$ (b) $\frac{1}{n}$
 (c) $\left(1 - \frac{1}{n^2}\right)$ (d) $\left(\frac{1}{n^2 - 1}\right)$

5. A particle of mass 4 m which is at rest explodes into three fragments. Two of the fragments each of mass m are found to move with a speed v each mutually perpendicular directions. The energy released in the process of explosion is -


- (a) $\frac{3}{2} mv^2$ (b) $3 mv^2$ (c) $2mv^2$ (d) $\frac{1}{2} mv^2$

6. Two pendulums start swinging together. Their lengths are respectively 1.44 m and 1 m . They will again start swinging together after -



- (a) 1 vibration (b) 3 vibrations
 (c) 4 vibrations (d) 5 vibrations

7. Gravitational field at the centre of a semicircle formed by a thin wire AB of mass m and length ℓ is -




- (a) $\frac{Gm}{\ell}$ along x-axis (b) $\frac{Gm}{\pi\ell}$ along y-axis
 (c) $\frac{2\pi Gm}{\ell^2}$ along x-axis (d) $\frac{2\pi GM}{\ell^2}$ along y-axis

8. A rod is non uniform having mass per unit length as μ which varies linearly with distance x as per relation $\mu = ax$ (a is a constant). If its total mass is M and length ℓ , the centre of mass is given by -

- (a) $\frac{3}{4} \ell$ (b) $\frac{2}{3} \ell$ (c) $\frac{2}{5} \ell$ (d) $\frac{\ell}{3}$

9. One end of a uniform rod of mass m_1 and cross sectional area A is hung from a ceiling. The other end of the bar is supporting mass m_2 . The stress at the midpoint is -




- (a) $\frac{g(m_2 + 2m_1)}{2A}$ (b) $\frac{g(m_2 + m_1)}{2A}$
(c) $\frac{g(2m_2 + m_1)}{2A}$ (d) $\frac{g(m_2 + m_1)}{A}$

10. A wave is represented by a equation $y = 0.5 \sin \frac{2\pi}{\lambda}(10t + x)$ metre. It is stated that it is a

travelling wave propagating along + x direction with velocity 10 ms^{-1} then it is determined that -

- (a) The statement is correct
(b) The statement is false
(c) The statement may be false or correct
(d) Nothing can be said because data is incomplete

11. Six moles an ideal gas performs a cycle show. If the temperatures are $T_A = 600\text{K}$, $T_B = 800\text{K}$, $T_C = 2200 \text{ K}$ and $T_D = 1200 \text{ K}$, the work done per cycle is -



- (a) 20000 J (b) 10000 J (c) 30000 J (d) 40000 J

12. Starting with the same initial conditions if an ideal gas expands from volume V_1 to V_2 in three different ways. The work done by gas is W_1 , if the process is purely isothermal, W_2 if purely isobaric and W_3 if purely adiabatic, then -

- (a) $W_2 > W_1 > W_3$ (b) $W_2 > W_3 > W_1$
(c) $W_1 > W_2 > W_3$ (d) $W_1 > W_3 > W_2$

13. Two capillaries of same length and radii in the ratio 1 : 2 are connected in series. A liquid flows through them in streamlined condition. If the pressure across the two extreme ends of the combination is 1 m of water, the pressure difference across first capillary is-


- (a) 9.4 m (b) 4.9 m (c) 0.49 m (d) 0.94 m

14. If the distance between the earth and the sun were half the present value then the number of days in a year could have been -
(a) 64.5 (b) 129 (c) 182.5 (d) 730

15. Potential difference between shell of an electrostatic generator and spray point is MV . If the transfer of charge to the shell is at the rate of Q unit per second, considering electrical forces only, the power provided to drive the belt is -


- (a) $\frac{VQ}{t} \times 10^6$ (b) $VQ \times 10^6$ (c) $\frac{VQ}{10^6}$ (d) $\frac{VQ}{t}$

16. Figure shows two equipotential lines in x, y plane for an electric field. Then x component $[E_x]$ and y component (E_y) of field in space between these lines are respectively -



- (a) $-100 \text{ Vm}^{-1}, 200 \text{ Vm}^{-1}$
(b) $-100 \text{ Vm}^{-1}, -200 \text{ Vm}^{-1}$
(c) $200 \text{ Vm}^{-1}, 100 \text{ Vm}^{-1}$
(d) $100 \text{ Vm}^{-1}, 100 \text{ Vm}^{-1}$

17. A dielectric slab is partially introduced between two square plates of area A of a parallel plates capacitor as shown. Dielectric constant of slab is ϵ_r . Total capacitance of the system is -



- (a) $\frac{\epsilon_0 \sqrt{Ax}}{d}$
(b) $\frac{\epsilon_0}{d} (A - \sqrt{Ax} + \epsilon_r \sqrt{Ax})$
(c) $\frac{\epsilon_0}{d} (\epsilon_r \sqrt{Ax} - A - \sqrt{Ax})$
(d) $\frac{\epsilon_0 \epsilon_r l}{d} (-\sqrt{Ax} + A + \epsilon_r \sqrt{Ax})$

18. A piece of copper and another of germanium are cooled from room temperature to 80 K. The resistance of -

- (a) Each of them increases
(b) Each of them decreases
(c) Cooper increases and germanium decreases
(d) Copper decreases and germanium increases


19. Two resistances $300\ \Omega$ and $400\ \Omega$ are connected in series with a battery of emf 60 V and negligible internal resistance. An ideal voltmeter reads 30 V when connected across $400\ \Omega$ resistor. The reading of same voltmeter across $300\ \Omega$ resistor is –

- (a) 19 V (b) 20 V (c) 22 V (d) 21 V

20. A potentiometer wire of length 100 cm has a resistance of $10\ \Omega$. It is connected with a resistance in series and an accumulator of emf 2 V and of negligible internal resistance. A source of emf 10 mV is balanced against a length of 40 cm of potentiometer wire. The value of external resistance is –


- (a) $790\ \Omega$ (b) $970\ \Omega$ (c) $97\ \Omega$ (d) $709\ \Omega$

21. The magnetic induction at point O. If the current carrying wire is in the shape shown –



- (a) $\frac{\mu_0 i}{4\pi r} \left[\frac{3}{2}\pi + 1 \right]$ (b) $\frac{\mu_0 i}{2\pi r} \left[\frac{3}{2}\pi + 1 \right]$
 (c) $\frac{\mu_0 i}{\pi r} \left[\frac{3}{2} \right]$ (d) $\frac{\mu_0 i}{2\pi r} \left[1 - \frac{3}{2}\pi \right]$

22. Some current I flows in two parallel conductors distant $2d$ as shown. The strength of magnetic field at a point P equidistant from both conductors is –




- (a) $\frac{\mu_0 Id}{4\pi a}$ (b) $\frac{\mu_0 Id}{2\pi a}$ (c) $\frac{\mu_0 Id}{4\pi a^2}$ (d) $\frac{\mu_0 Id}{\pi a^2}$

23. A coil in the shape of an equilateral triangle of side 0.02 m is suspended from a vertex such that it is hanging in a vertical plane between the pole pieces of a permanent magnet producing a horizontal magnetic field of $5 \times 10^{-2}\text{ T}$. The couple acting on coil when a current of 0.1 A is passed through it and the magnetic field is parallel to its plane –

- (a) $6.86 \times 10^{-7}\text{ Nm}$ (b) $8.66 \times 10^{-7}\text{ Nm}$
 (c) $8.7 \times 10^{-9}\text{ Nm}$ (d) $8 \times 10^{-8}\text{ Nm}$

24. A Thin prism of angle A and refractive index μ for sodium light is placed at a distance S from a slit illuminated by sodium light. Distance between slit and image formed by prism is –



- (a) $As(1 - \mu)$ (b) $As(1 + \mu)$
 (c) $As(\mu - 1)$ (d) $\frac{A}{S}(1 - \mu)$

25. The resolution limit of eye is 60 s . A distance of $x\text{ km}$ from the eye two persons stand with lateral separation of 3 m . For the two persons to be just resolved by eye. x should be –

- (a) 10 km (b) 15 km (c) 20 cm (d) 30 km

26. Yellow light is used in a single slit diffraction experiment with slit width of a 0.6 mm . If yellow light is replaced by x-rays then the observed pattern will reveal –

- (a) That the central maximum is narrower
 (b) more number of fringes
 (c) less number of fringes
 (d) no diffraction patterns

27. Two beams of light having intensities I and $4I$ interfere to produce a fringe pattern on a screen. The phase difference between the beams is $\frac{\pi}{2}$ at point A and π at point B. Then

the difference between the resultant intensities at A and B is–

- (a) $2I$ (b) $4I$ (c) $5I$ (d) $7I$

28. A proton and an α particle have KE in the ratio of $16 : 1$. The ratio of de-Broglie waves associated with them is –

- (a) $1 : 2$ (b) $2 : 1$ (c) $2 : 3$ (d) $1 : 4$






29. Light of wavelength $5000\ \text{\AA}$ falls on a sensitive surface. If the surface has received 10^{-7} J of energy then number of photons just falling on the surface are–





- (a) 2.5×10^6 (b) 2.5×10^{11}
 (c) 2.5×10^3 (d) 5000

30. The wavelength of K_α line produced by an x-ray tube of material is $0.76\ \text{\AA}$. The atomic number of material of anode tube is –

- (a) 41 (b) 14 (c) 51 (d) 15

CHEMISTRY

31. 1.25g of a solid dibasic acid is completely neutralized by 25 ml. of 0.25 molar Ba(OH)₂ solution. Molecular mass of the acid is -
 (a) 100 (b) 150 (c) 120 (d) 200
32. Rates of effusion of hydrogen and deuterium under similar conditions are in the ratio -
 (a) 1 : 1 (b) $\sqrt{2}$: 1 (c) 2 : 1 (d) 1 : 4
33. For equilibrium
 $\text{NH}_4\text{HS(s)} \rightleftharpoons \text{NH}_3\text{(g)} + \text{H}_2\text{S(g)}$
 $K_c = 1.8 \times 10^{-4}$ at 298 K.
 The value of K_p at 298 K is -
 (a) 0.108 (b) 4.4×10^{-3}
 (c) 1.8×10^{-4} (d) 4.4×10^{-4}
34. Given that
 $\text{H}_2\text{O(l)} \rightarrow \text{H}_2\text{O(g)} ; \Delta H = + 43.7 \text{ kJ}$
 $\text{H}_2\text{O(s)} \rightarrow \text{H}_2\text{O(l)} ; \Delta H = + 6.05 \text{ kJ}$
 $\Delta H_{\text{sublimation of ice}}$ is -
 (a) 49.75 kJ mol⁻¹ (b) 37.65 kJ mol⁻¹
 (c) 43.7 kJ mol⁻¹ (d) - 43.67 kJ mol⁻¹
35. The solubility product K_{sp} of sparingly soluble salt Ag₂CrO₄ is 4×10^{-12} . The solubility of the salt is -
 (a) $1 \times 10^{-12} \text{ M}$ (b) $2 \times 10^{-6} \text{ M}$
 (c) $1 \times 10^{-6} \text{ M}$ (d) $1 \times 10^{-4} \text{ M}$
36. Which of the following chemical reactions depicts the oxidising behaviour of H₂SO₄ ?
 (a) $2\text{HI} + \text{H}_2\text{SO}_4 \rightarrow \text{I}_2 + \text{SO}_2 + 2\text{H}_2\text{O}$
 (b) $\text{Ca(OH)}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + 2\text{H}_2\text{O}$
 (c) $\text{NaCl} + \text{H}_2\text{SO}_4 \rightarrow \text{NaHSO}_4 + \text{HCl}$
 (d) $2\text{PCl}_5 + \text{H}_2\text{SO}_4 \rightarrow 2\text{POCl}_3 + 2\text{HCl} + \text{SO}_2\text{Cl}_2$
37. Potassium has a bcc structure with nearest neighbour distance of 4.52 Å. If atomic mass of potassium is 3a, its density is -
 (a) 454 kg m⁻³ (b) 804 kg m⁻³
 (c) 852 kg m⁻³ (d) 900 kg m⁻³
38. If $E_{\text{Zn}^{2+}/\text{Zn}}^0 = -0.763 \text{ V}$
 and $E_{\text{Cd}^{2+}/\text{Cd}}^0 = -0.403 \text{ V}$, the emf of the cell
 $\text{Zn} | \text{Zn}^{2+} || \text{Cd}^{2+} | \text{Cd}$ ($a = 0.004$), ($a = 0.2$) will be given by -
 (a) $E = -0.36 + \frac{0.059}{2} \log \frac{0.004}{2}$
 (b) $E = +0.36 + \frac{0.059}{2} \log \frac{0.04}{2}$
 (c) $E = -0.36 + \frac{0.059}{2} \log \frac{0.2}{0.004}$
 (d) $E = +0.36 + \frac{0.059}{2} \log \frac{0.2}{0.004}$
39. The value of P° for benzene of certain temperature is 640 mm of Hg. The vapour pressure of solution containing 2.5 g of a certain substance 'A' in 39.0 g of benzene is 600 mm of Hg. The molecular mass of A is -
 (a) 65.25 (b) 130 (c) 40 (d) 80
40. A reaction which is of first order w.r.t. reactant A, has a rate constant 6 min⁻¹. If we start with $[\text{A}] = 0.5 \text{ mol L}^{-1}$, when would $[\text{A}]$ reach the value of 0.05 mol L⁻¹ ?
 (a) 0.384 min (b) 0.15 min
 (c) 3 min (d) 3.84 min
41. $\text{CH}_3\text{NH}_2 + \text{CHCl}_3 + \text{KOH} \rightarrow$ Nitrogen containing compound + KCl + H₂O
 Nitrogen containing compound is -
 (a) $\text{CH}_3 - \text{C} \equiv \text{N}$ (b) $\text{CH}_3 - \text{NH} - \text{CH}_3$
 (c) $\text{CH}_3 - \text{N} \equiv \text{C}^+$ (d) $\text{CH}_3 - \text{N}^+ \equiv \text{C}^-$
42. 4-methyl benzene sulphonic acid react with sodium acetate to give -

 (a)  ; CH₃COOH (b)  ; SO₃
 (c)  ; SO₃ (d)  ; NaOH
43. The product(s) obtained via oxymercuration ($\text{HgSO}_4 + \text{H}_2\text{SO}_4$) of 1-butyne would be -
 (a) $\text{CH}_3\text{CH}_2\text{COCH}_3$
 (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$
 (c) $\text{CH}_3\text{CH}_2\text{CHO} + \text{HCHO}$
 (d) $\text{CH}_3\text{CH}_2\text{COOH} + \text{HCOOH}$
44. Acetophenone is prepared by the reaction of which of the following in the presence of AlCl₃ catalyst -
 (a) Phenol and acetic acid
 (b) Benzene and acetone
 (c) Benzene and acetyl chloride
 (d) Phenol and acetone




where M = molecule


R = Reagent

M and R are


- (a) CH_3CH_2Cl and $NaOH$
 (b) $CH_2Cl - CH_2OH$ and aq. $NaHCO_3$
 (c) CH_3CH_2OH and HCl
 (d) $CH_2 = CH_2$ and heat



50. Which is least reactive towards nucleophilic substitution (SN_2)




51. Among the following the least stable resonance structure is –



52. Homolytic fission of C–C bond in ethane gives an intermediate in which carbon is –

- (a) sp^3 hybridised (b) sp^2 hybridised
 (c) sp hybridised (d) sp^2d hybridised

53. The IUPAC name of the compound  is–

- (a) (2E, 4E)-2, 4-hexadiene
 (b) (2Z, 4Z)-2, 4-hexadiene
 (c) (2Z, 4E)-2, 4-hexadiene
 (d) (2E, 4Z)-4, 2-hexadiene

54. The brown ring test for NO_2^- and NO_3^- is due to the formation of complex ion with the formula –

- (a) $[Fe(H_2O)_6]^{2+}$ (b) $[Fe(NO)(CN)_5]^{2+}$
 (c) $[Fe(H_2O)_5NO]^{2+}$ (d) $[Fe(H_2O)(NO)_5]^{2+}$

55. The correct order for the wavelength of absorption in the visible region is –

- (a) $[Ni(NO_2)_6]^{4-} < [Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+}$
 (b) $[Ni(NO_2)_6]^{4-} < [Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+}$
 (c) $[Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
 (d) $[Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+} < [Ni(NO_2)_6]^{4-}$

56. In nitroprusside ion, the iron and NO exists as Fe (II) and NO^+ rather than Fe(III) and NO these forms can be differentiated by –

- (a) Estimating the concentration of iron
 (b) Measuring the concentration of CN^-
 (c) Measuring the solid state magnetic moment
 (d) Thermally decomposing the compound

57. Four reactions are given below
 I $2\text{Li} + 2\text{H}_2\text{O} \rightarrow 2\text{LiOH} + \text{H}_2$
 II $2\text{Na} + 2\text{H}_2\text{O} \rightarrow 2\text{NaOH} + \text{H}_2$
 III $2\text{LiNO}_3 \xrightarrow{\text{heat}} 2\text{LiNO}_2 + \text{O}_2$
 IV $2\text{NaNO}_3 \xrightarrow{\text{heat}} 2\text{NaNO}_2 + \text{O}_2$
 Which of the above if any is wrong
 (a) IV (b) III (c) I (d) None
58. Name of the structure of silicates in which three oxygen atoms of $[\text{SiO}_4]^{4-}$ are shared is –
 (a) Pyrosilicate
 (b) Sheet silicate
 (c) Linear chain silicate
 (d) Three dimensional silicate
59. 'Lapis-Lazuli' is a blue coloured precious stone. It is mineral of the class –
 (a) Sodium aluminosilicate
 (b) Zinc-cobaltate
 (c) Basic copper carbonate
 (d) Prussian blue
60. In which of the following arrangements the order is not according to the property indicating against it –
 (a) $\text{Al}^{3+} < \text{Mg}^{2+} < \text{Na}^+ < \text{F}^-$ (increasing ionic size)
 (b) $\text{B} < \text{C} < \text{N} < \text{O}$ (increasing first I.E.)
 (c) $\text{I} < \text{Br} < \text{F} < \text{Cl}$
 (increasing electron gain enthalpy (–ve))
 (d) $\text{Li} < \text{Na} < \text{K} < \text{Rb}$ (increasing metallic radius)
64. The relation R defined on the set $A = \{1, 2, 3\}$ is given by $R = \{(1, 1), (2, 2)\}$ then number of correct choices from the following is –
 (i) reflexive (ii) symmetric
 (iii) Transitive (iv) anti symmetric
 (a) 1 (b) 2 (c) 3 (d) 4
65. Let U be the universal set and $A \cup B \cup C = U$ then $\{(A - B) \cup (B - C) \cup (C - A)\}^c =$
 (a) $A \cap (B \cap C)$ (b) $A \cap (B \cup C)$
 (c) $(A \cap B \cap C)$ (d) None of these
66. If A and B are square matrices of same size and $|B| \neq 0$ then $(B^{-1}AB)^4 =$
 (a) $(B^4)^{-1}AB^4$ (b) BA^4B^{-1}
 (c) $B^{-1}A^4B$ (d) None of these
67. Let $g(x) = \begin{vmatrix} f(x+\alpha) & f(x+2\alpha) & f(x+3\alpha) \\ f(\alpha) & f(2\alpha) & f(3\alpha) \\ f'(\alpha) & f'(2\alpha) & f'(3\alpha) \end{vmatrix}$
 where α is a constant then $\lim_{x \rightarrow 0} \frac{g(x)}{x} =$
 (a) 0 (b) 1 (c) –1 (d) None
68. If $\Delta_1 = \begin{vmatrix} f & 2d & e \\ 2z & 4x & 2y \\ e & 2a & b \end{vmatrix}$ and $\Delta_2 = \begin{vmatrix} 2a & b & e \\ 2d & e & f \\ 4x & 2y & 2z \end{vmatrix}$
 then $\Delta_1/\Delta_2 =$
 (a) 1 (b) 2 (c) $\frac{1}{2}$ (d) None

MATHEMATICS

61. The mean of 0, 1, 2, 3, ..., n, with the corresponding weight ${}^nC_0, {}^nC_1, {}^nC_2, \dots, {}^nC_n$ is
 (a) $\frac{2^n}{n+1}$ (b) $\frac{2^{n+1}}{n(n+1)}$ (c) $\frac{n+1}{2}$ (d) $\frac{n}{2}$
62. The function $f(x) = 1 + x \sin x [\cos x]$,
 $0 < x \leq \frac{\pi}{2}$ where $[\cdot] = \text{G.I.F.}$
 (a) is continuous on $(0, \pi/2)$
 (b) is strictly decreasing in $(0, \pi/2)$
 (c) is strictly increasing in $(0, \pi/2)$
 (d) has global maximum value 2
63. If the radius of a spherical balloon is measured with in 1%, the error (in percent) in the volume is –
 (a) $4\pi r^2\%$ (b) 3% (c) $\left(\frac{88}{7}\right)\%$ (d) None
69. The number of ways in which 20 one rupee coin can be distributed among 5 people such that each person gets at least 3 rupee is –
 (a) 26 (b) 63 (c) 125 (d) None
70. The total number of six digit number $x_1 x_2 x_3 x_4 x_5 x_6$ have the property that $x_1 < x_2 \leq x_3 < x_4 < x_5 \leq x_6$ is equal to –
 (a) ${}^{10}C_6$ (b) ${}^{12}C_6$ (c) ${}^{11}C_6$ (d) None
71. $2 \left\{ 1 + \frac{a^2}{\lfloor 2 \rfloor} + \frac{a^4}{\lfloor 4 \rfloor} + \dots \right\}$; $a = \log_e n$ is equal to
 (a) $\frac{n-1}{n}$ (b) $\frac{n^2-1}{n}$ (c) $\frac{n+1}{n}$ (d) $\frac{n^2+1}{n}$
72. The term independent of x in $\left(\sqrt{x} - \frac{2}{x}\right)^{18}$ is –
 (a) ${}^{18}C_6 2^6$ (b) ${}^{18}C_{12} 2^{12}$ (c) ${}^{18}C_8 2^8$ (d) None

73. $1 + \frac{2}{3} \cdot \frac{1}{2} + \frac{2.5}{3.5} \left(\frac{1}{2}\right)^2 + \frac{2.5.8}{3.6.9} \left(\frac{1}{2}\right)^3 + \dots =$
 (a) $2^{1/3}$ (b) $3^{1/4}$ (c) $4^{1/3}$ (d) $3^{1/3}$
74. If ω is imaginary cube root of unity then $\arg(i\omega) + \arg(i\omega^2) =$
 (a) 0 (b) $\pi/2$ (c) π (d) None
75. $\sum_{r=1}^n \frac{1}{\log_2 4^r}$ is equal to
 (a) $\frac{n(n+1)}{4}$ (b) $\frac{n(n+1)}{2}$ (c) $n(n+1)$ (d) None
76. If a_1, a_2, \dots, a_{15} are in A.P. and $a_1 + a_8 + a_{15} = 15$ then $a_2 + a_3 + a_8 + a_{13} + a_{14} =$
 (a) 15 (b) 10 (c) 25 (d) None
77. If $a_1 < a_2 < a_3 < a_4 < a_5 < a_6$ then the equation $(x - a_1)(x - a_3)(x - a_5) + 2(x - a_2)(x - a_4)(x - a_6) = 0$ has
 (a) Four real roots
 (b) One real root
 (c) One real root in each interval $(a_1, a_2), (a_3, a_4)$ and (a_5, a_6)
 (d) None of these
78. Solution of the differential equation $x dx + z dy + (y + 2z) dz = 0$ is
 (a) $x^2 + 2yz + 2z^2 = c$ (b) $x^2 + yz + z^2 = c$
 (c) $x^2 + 2yz + z^2 = c$ (d) None of these
79. The slope of the tangent to the curve $y = f(x)$ at $(x, f(x))$ is $(2x + 1)$. If the curve passes through the point $(1, 2)$, then the area bounded by the curve, x-axis and the lines $x = 1, x = 0$ is
 (a) $5/6$ (b) $6/5$ (c) 6 (d) 1
80. The maximum area of a rectangle whose two consecutive vertices lie on the x-axis and another two lie on the curve $y = e^{-|x|}$ is equal to
 (a) $2e$ (b) $2/e$ (c) e (d) $1/e$
81. $\int \sqrt{\sin^2 x} dx =$
 (a) $-\cos x + C$ (b) $\cos x + C$
 (c) $-\cos x \operatorname{sgn} \sin x + C$ (d) None of these
82. The maximum value of $P = 6x + 8y$ subject to constrains $2x + y \leq 30, x + 2y \leq 24$ and $x \geq 0, y \geq 0$ is -
 (a) 90 (b) 120 (c) 96 (d) 240
83. Segment of the tangent to the curve $xy = c^2$ at the point (x', y') which is contained between the co-ordinate axes is bisected at the point
 (a) $(-x', y')$ (b) (y', x') (c) $\left(\frac{x'}{2}, \frac{y'}{2}\right)$ (d) None
84. There is a point $P(a, a, a)$ on the line passing through the origin and equally inclined with axes the equation of the plane perpendicular to OP and passing through P cuts the intercepts on axes the sum of whose reciprocals is
 (a) a (b) $3/2a$ (c) $3a/2$ (d) $1/a$
85. If $\vec{a} = p\hat{i} + 5\hat{j} + 17\hat{k}$ and $\vec{b} = 2\sqrt{q}\hat{i} + 13\hat{j} + \hat{k}$ have equal magnitude and p, q are positive integer $\in [1, 1000]$ then the total number of ordered pair (p, q) is
 (a) 33 (b) 32 (c) 31 (d) None
86. The equation $\frac{x^2}{8-a} + \frac{y^2}{a-2} = 1$ will represent an ellipse if
 (a) $a \in (1, 4)$ (b) $a \in (-\infty, 2) \cup (8, \infty)$
 (c) $a \in (2, 8)$ (d) None of these
87. Angle between the tangent drawn to $y^2 = 4x$ at the point where it is intersected by line $y = x - 1$ is
 (a) $\pi/6$ (b) $\pi/3$ (c) $\pi/4$ (d) $\pi/2$
88. Consider four circles $(x \pm 1)^2 + (y \pm 1)^2 = 1$. Equation of smaller circle touching these four circles is
 (a) $x^2 + y^2 = 3 - \sqrt{2}$ (b) $x^2 + y^2 = 6 - 3\sqrt{2}$
 (c) $x^2 + y^2 = 5 - 2\sqrt{2}$ (d) $x^2 + y^2 = 3 - 2\sqrt{2}$
89. If the point $P(a, a^2)$ lies completely inside the triangle formed by the lines $x = 0, y = 0$ and $x + y = 2$ then exhaustive range of 'a' is
 (a) $a \in (0, 1)$ (b) $a \in (1, \sqrt{2})$
 (c) $a \in (\sqrt{2} - 1, \sqrt{2})$ (d) $a \in (\sqrt{2} - 1, 1)$
90. If the sides of a Δ are $3 : 7 : 8$ then $R : r$ is equal to
 (a) $2 : 7$ (b) $7 : 2$ (c) $3 : 7$ (d) None
91. The function $f(x) = \frac{x}{1 + x \tan x}, \left(0, \frac{\pi}{2}\right)$ has
 (a) One point of minimum
 (b) One point of maximum
 (c) No extreme point
 (d) Two point of maximum

92. If solution of the equation $3\cos^2\theta - 2\sqrt{3}\sin\theta\cos\theta - 3\sin^2\theta = 0$ are $n\pi + \frac{\pi}{r}$ and $n\pi + \frac{\pi}{s}$ then $|r - s| =$
 (a) 3 (b) 9 (c) 7 (d) 1
93. If $\cot^{-1} \frac{n}{\pi} > \frac{\pi}{6}$, $n \in \mathbb{N}$ then maximum value of n is
 (a) 6 (b) 5 (c) 4 (d) 3
94. Period of the function $f(x) = \sin 3\pi\{x\} + \tan \pi[x]$ where $[.]$ and $\{.\}$ denote the integral part and fractional part respectively, is given by
 (a) 1 (b) 2 (c) 3 (d) π
95. The domain and range of $f(x) = \cos^{-1} \sqrt{\log_{[x]} \left(\frac{|x|}{x} \right)}$ where $[.]$ denotes the greatest integer function respectively
 (a) $[1, \infty)$, $\left[0, \frac{\pi}{2}\right]$ (b) $[2, \infty)$, $\left[0, \frac{\pi}{2}\right]$
 (c) $[2, \infty)$, $\left\{\frac{\pi}{2}\right\}$ (d) $[1, \infty)$, $\{0\}$
96. The graph of the function $y = f(x)$ has a unique tangent not parallel to x -axis at the point $(a, 0)$ through which the graph passes, then $\lim_{x \rightarrow a} \frac{\log_e \{1 + 6f(x)\}}{3f(x)}$ is
 (a) 1 (b) 0 (c) 2 (d) None
97. If $P = \lim_{x \rightarrow 5^+} \frac{x^2 - 9x + 20}{x - [x]} - \lim_{x \rightarrow 4^-} \frac{x^2 - 9x + 20}{x - [x]}$
 and $Q = \lim_{x \rightarrow 4^+} \frac{x^2 - 9x + 20}{x - [x]} - \lim_{x \rightarrow 5^-} \frac{x^2 - 9x + 20}{x - [x]}$
 where $[.] = \text{G.I.F.}$ then $\frac{P}{Q} =$
 (a) 1 (b) 2 (c) 3 (d) None
98. If $(a + bx)e^{y/x} = x$ then $\frac{1}{y_2} (xy_1 - y)^2 =$
 (A) x^3 (b) $3x^2$ (c) $1/x^3$ (d) None
99. If $f(x)$ is continuous function such that $\int_n^{n+1} f(x) dx = n^3$, $n \in \mathbb{Z}$ then $\int_{-2}^3 f(x) dx =$
 (a) 16 (b) 0 (c) 2 (d) None

100. If $x^{2f(x)} + f\left(\frac{1}{x}\right) = 2$ for all x except at $x = 0$
 then $\int_{1/3}^3 f(x) dx =$
 (a) $4/3$ (b) $8/3$ (c) $1/3$ (d) None

ENGLISH PROFICIENCY

101. Find the correctly spelt word –
 (a) Geraff (b) Giraffe (c) Giraf (d) Gerraffe
102. Pick up the correct synonym for the following words
Plush :
 (a) Luxurious (b) Delicious
 (c) Comforting (d) Tasty
103. Choose the alternative which can replace the word printed in underline without changing the meaning of the sentence.
 When he returned, he was accompanied by 'sprightly' young girl.
 (a) Lively (b) Beautiful
 (c) Sportive (d) Intelligent
104. Choose the word which is closest to the 'opposite' in meaning of the underlined word
 Many snakes are 'innocuous' :
 (a) Deadly (b) Ferocious
 (c) Poisonous (d) Harmful
105. Choose the one which can be substituted for the given words/sentences :
'Giving undue favours to one's kith and kin'
 (a) Corruption (b) Worldliness
 (c) Favouritism (d) Nepotism
106. Find out which one of the words given below the sentence can most appropriately replace the group of words underlined in the sentence :
 The bus has to "go back and forth" every six hours.
 (a) Cross (b) Shuttle (c) Travel (d) Run
107. Which one of the two sentences given below is wrong on the basis of the underlined words :
 1. He is a very "ingenuous" business-man.
 2. I like him for his "ingenious" nature.
 (a) Sentence 1 is correct
 (b) Sentence 2 is correct
 (c) Both the sentences can be made correct by interchanging the underlined words.
 (d) Both the sentences can not be interchanged hence, both are wrong

108. Choose from the given words below the two sentences, that word which has the same meaning and can be used in the same context as the part given underlined in both the sentences :

1. His "aloof" behaviour is an indication of his arrogance.
 2. During our field visits we visited "remote" parts of Rajasthan.
- (a) Far-off (b) Introvert
(c) Distant (d) Depressed

109. Find out which part of the sentence has an error. If there is no mistake, the answer is 'No error'.

- "Meatlessdays" / havebeen made / into a film / No Error
(a) (b) (c) (d)
- (a) Meatless days (b) have been made
(c) into a film (d) No Error

110. Replace the underlined word with one of the given options :

- The Second World War started in 1939.
- (a) Broke out (b) Set out
(c) Took out (d) Went out

LOGICAL REASONING

111. Fill in the blank space

- 6, 13, 28, . ? . .
(a) 56 (b) 57 (c) 58 (d) 59


112. Choose the best alternative

- Car : Petrol :: T.V. : ?
(a) Electricity (b) Transmission
(c) Entertainment (d) Antenna

113. Pick the odd one out


- (a) Titan (b) Mercury
(c) Earth (d) Jupiter

114. **Direction** : In questions, find out which of the figures (a), (b), (c) and (d) can be formed from the pieces given in (x).




115. **Direction** : In question, choose the set of figures which follows the given rule.


Rule : Closed figures become more and more open and open figures more and more closed.




116. **Direction** : In question below, you are given a figure (x) followed by four figures (a), (b), (c) and (d) such that (x) is embedded in one of them. Trace out the correct alternative.




117. **Direction** : In following question, find out which of the answer figures (a), (b), (c) and (d) completes the figure-matrix ?




118. **Direction** : The questions that follow contain a set of three figure X, Y and Z showing a sequence of folding of piece of paper. Fig. (Z) shows the manner in which the folded paper has been cut. These three figure are followed by four answer figure from which you have to choose a figure which would most closely resemble the unfolded form of figure. (Z)




119.Direction : In following question, complete the missing portion of the given pattern by selecting from the given alternatives (a), (b), (c) and (d).




(X)



120.Direction : In question below, you are given a figure (X) followed by four figures (a), (b), (c) and (d) such that (X) is embedded in one of them. Trace out the correct alternative.



(X)



121. In a certain code, ROAD is written as URDG. How is SWAN written in that code ?

- (a) VXDQ (b) VZDQ (c) VZCP (d) UXDQ

122. If ZIP = 198 and ZAP = 246, then how will you code VIP ?

- (a) 174 (b) 222 (c) 888 (d) 990

123. Select the related letter/word/number from the given alternatives

Train : Passenger :: Aeroplane : ?

- (a) Aeronaut (b) Astronaut
(c) Pilot (d) Air hostess

124. Four words have been given, out of which three are alike in some manner and the fourth one is different. Choose out the odd one.

- (a) Wood (b) Cork (c) Stone (d) Paper

125. Choose the correct alternatives from the given ones that will complete the series.

2, 12, 30, 2, 90, 132

- (a) 48 (b) 56 (c) 63 (d) 72


126. Choose the correct alternatives from the given ones that will complete the series.

a _ b _ ba _ b _ _ ba


- (a) bbaab (b) bbabb (c) aabab (d) aabba

127. Which of the following cube in the answer figure be made based on the unfolded cube in the question figure ?

Question Figure




Answer figure




- (a) Only (a) (b) Only (a) and (d)
(c) Only (b) and (c) (d) Only (a) and (d)

128. Which figure represent the relationship among

Sun, Moon, Molecule?




129. Select the missing number from the given responses.




- (a) 189 (b) 227 (c) 277 (d) 339

130. Identify the figure that will complete the pattern.

Question Figure :



Answer Figure :



ANSWERS

PHYSICS

- 1.(b) 2.(a) 3.(d) 4.(c) 5.(a) 6.(d) 7.(d) 8.(b) 9.(c) 10.(b)
 11.(d) 12.(a) 13.(d) 14.(b) 15.(b) 16.(a) 17.(b) 18.(d) 19.(d) 20.(a)
 21.(a) 22.(d) 23.(b) 24.(c) 25.(a) 26.(d) 27.(b) 28.(a) 29.(b) 30.(a)

CHEMISTRY

- 31.(d) 32.(b) 33.(a) 34.(a) 35.(d) 36.(a) 37.(d) 38.(b) 39.(d) 40.(a)
 41.(d) 42.(a) 43.(a) 44.(c) 45.(b) 46.(a) 47.(a) 48.(b) 49.(a) 50.(c)
 51.(a) 52.(b) 53.(a) 54.(c) 55.(a) 56.(c) 57.(b) 58.(d) 59.(a) 60.(b)

MATHEMATICS

- 61.(d) 62.(a) 63.(b) 64.(c) 65.(c) 66.(c) 67.(a) 68.(a) 69.(d) 70.(c)
 71.(d) 72.(a) 73.(c) 74.(d) 75.(a) 76.(c) 77.(c) 78.(a) 79.(a) 80.(b)
 81.(c) 82.(b) 83.(d) 84.(d) 85.(c) 86.(d) 87.(d) 88.(d) 89.(a) 90.(b)
 91.(b) 92.(b) 93.(b) 94.(a) 95.(c) 96.(c) 97.(d) 98.(a) 99.(b) 100.(b)

ENGLISH PROFICIENCY

- 101.(b) 102.(a) 103.(a) 104.(d) 105.(d) 106.(b) 107.(c) 108.(c) 109.(a) 110.(a)

LOGICAL REASONING

- 111.(d) 112.(a) 113.(a) 114.(c) 115.(b) 116.(d) 117.(b) 118.(a) 119.(c) 120.(a)
 121.(b) 122.(b) 123.(a) 124.(c) 125.(b) 126.(b) 127.(a) 128.(c) 129.(b) 130.(b)

Hints & Solutions

PHYSICS

1. [b] $V_2 = V_1 \frac{\alpha^2}{\beta}$

i.e. $[L_2 T_2^{-1}] = [L_1 T_1^{-1}] \frac{\alpha^2}{\beta}$... (i)

$a_2 = a_1 \alpha \beta$

i.e. $[L_2 T_2^{-2}] = [L_1 T_1^{-2}] \alpha \beta$... (ii)

Also $F_2 = \frac{F_1}{\alpha \beta}$

i.e. $[M_2 L_2 T_2^{-2}] = [M_1 L_1 T_1^{-2}] \frac{1}{\alpha \beta}$... (iii)

Dividing equation (iii) by equation (ii)

$$M_2 = \frac{M_1}{(\alpha \beta)(\alpha \beta)} = \frac{M_1}{\alpha^2 \beta^2}$$

Squaring equation (i) and dividing by equation (ii)

$$\frac{[L_2 T_2 T_2^2]}{[L_2 T_2^{-2}]} = \frac{[L_1^2 T_1^{-2}]}{[L_1 T_1^{-2}] \alpha \beta} \frac{\alpha^4}{\beta^2} \text{ or } L_2 = L_1 \frac{\alpha^3}{\beta^3}$$

Dividing equation (i) by equation (ii)

$$\frac{1}{T_2^{-1}} = \frac{1}{T_1^{-2}} \frac{\alpha^2}{\beta \alpha \beta} \text{ or } T_2 = T_1 \frac{\alpha}{\beta^2}$$

2. [a] Using $H = \frac{u^2 \sin \alpha}{2g}$ and

$$R = \frac{2u^2 \sin \alpha \cos \alpha}{g}$$

We get, $R^2 = \frac{4u^2}{g^2} \sin^2 \alpha \cos^2 \alpha$

Eliminating α ,

$$R^2 = \frac{4u^2}{g} \frac{2gH}{u^2} \left(1 - \frac{2gH}{u^2}\right) = \frac{8H}{g} (u^2 - 2gH)$$

$$\begin{aligned} \text{or } R &= \left[\frac{8H}{g} (u^2 - 2gH) \right]^{1/2} \\ &= \left[\frac{8 \times 25}{9.8} (40^2 - 2 \times 9.8 \times 25) \right]^{1/2} \\ &= 150.5 \text{ m} \end{aligned}$$

3. [d] Let $P = Q = x$ and $R = \sqrt{2x}$

using $\vec{P} + \vec{Q} + \vec{R} = 0$,

we get $\vec{P} + \vec{Q} = -\vec{R}$

$$(\vec{P} + \vec{Q}) \cdot (\vec{P} + \vec{Q}) = (-\vec{R}) \cdot (-\vec{R})$$

Then $P^2 + Q^2 + 2PQ \cos \theta = R^2$

i.e., $x^2 + x^2 + 2x^2 \cos \theta = 2x^2$

i.e. $\cos \theta = 0$, or $\theta = 90^\circ$

Again $\vec{Q} + \vec{R} = -\vec{P}$

$$(\vec{Q} + \vec{R}) \cdot (\vec{Q} + \vec{R}) = (-\vec{P}) \cdot (-\vec{P})$$

Then $Q^2 + R^2 + 2QR \cos \alpha = P^2$

i.e., $x^2 + 2x^2 + 2\sqrt{2} x^2 \cos \alpha = x^2$

i.e. $\cos \alpha = -\frac{1}{\sqrt{2}}$ or $\phi = 135^\circ$

Third angle = $360 - (135 + 90) = 135^\circ$

4. [c] Let a be the acceleration down the rough plane and a' be the acceleration down the frictionless plane. Taking L as the length of the inclined plane, we get

$$a = g(\sin \theta - \mu \cos \theta)$$

$$= g \left(\frac{1}{\sqrt{2}} - \frac{\mu}{\sqrt{2}} \right) \quad (\because \theta = 45^\circ)$$

and $a' = g \sin \theta = g \frac{1}{\sqrt{2}}$

Then, $L = \frac{1}{2} a t_1^2 = \frac{1}{2} a' t_2^2$


$$\text{or } \frac{1}{2} g \left(\frac{1}{\sqrt{2}} - \frac{\mu}{\sqrt{2}} \right) t^2 = \frac{1}{2} \frac{g}{\sqrt{2}} t_2^2$$

But $t_1 = n t_2 \dots$ (given)

$$\therefore \frac{1}{2} g \left(\frac{1}{\sqrt{2}} - \frac{\mu}{\sqrt{2}} \right) n^2 t_2^2 = \frac{1}{2} \frac{g}{\sqrt{2}} t_2^2$$

$$\text{or } 1 = (1 - \mu)n^2 \quad \text{or } \mu = \left(1 - \frac{1}{n^2}\right)$$

5. [a] Here momentum of third fragment is



$$P_3 = \sqrt{P_1^2 + P_2^2}$$

$$\text{or } P_3 = \sqrt{(mv)^2 + (mv)^2} = \sqrt{2} mv$$

Final KE of the system

$$= \frac{P_1^2}{2m} + \frac{P_2^2}{2m} + \frac{P_3^2}{2(2m)}$$

$$= \frac{1}{2} mv^2 + \frac{1}{2} mv^2 + \frac{1}{2} mv^2 = \frac{3}{2} mv^2$$

Since initial KE = 0 therefore energy released

$$= \frac{3}{2} mv^2$$


6. [d] Here $nT_1 = (n+1)T_2$

$$\text{i.e. } \frac{n}{n+1} = \frac{T_2}{T_1} = \sqrt{\frac{\ell_2}{\ell_1}} \quad \left[\because T = 2\pi \sqrt{\frac{\ell}{g}} \right]$$

$$= \sqrt{\left(\frac{1}{1.44}\right)} = \frac{10}{12}$$

or $2n = 10$ i.e. $n = 5$ vibrations

7. [d] Let a unit mass be kept at O then taking $d\ell = r d\theta$ as a small element and m/ℓ as mass per unit length, we get



$$d\vec{E} = \frac{GM}{\ell} \frac{rd\theta}{r^2} (\hat{i} \cos \theta d\theta + \hat{j} \sin \theta d\theta)$$

$$\int_0^\pi d\vec{E} = \int_0^\pi \frac{GM}{\ell} \frac{rd\theta}{r^2} (\hat{i} \cos \theta d\theta + \hat{j} \sin \theta d\theta)$$

$$\text{or } \vec{E} = \frac{2\pi GM}{\ell^2} \text{ along vertical (y axis) direction}$$

8. [b] A rod is non uniform having mass per unit length as Here

$$X_{cm} = \frac{1}{M} \int_0^{\ell} ndm = \frac{1}{M} \int_0^{\ell} x(\mu dx)$$

$$= \frac{1}{M} \int_0^{\ell} x(ax)dx = \frac{a}{M} \int_0^{\ell} x^2 dx = \frac{a\ell^3}{3M}$$

$$\text{Again } M = \int_0^{\ell} \mu dx = \int_0^{\ell} ax dx = \frac{a\ell^2}{2}$$

$$\therefore X_{cm} = \frac{a\ell^3 \times 2}{3(a\ell^2)} = \frac{2}{3} \ell$$

9. [c] Using stress = $\frac{F}{A}$, we get

$$\text{Stress at midpoint} = \frac{(m_2 g + \frac{m_1}{2} g)}{A}$$

$$= \frac{g(2m_2 + m_1)}{2A}$$

10. [b] Comparing the given equation with

$$y = A \sin \frac{2\pi}{\lambda} (vt + x)$$

We find the $v = 10 \text{ ms}^{-1}$ but the wave is travelling along -ve x-axis because there is +ve sign between vt and x.

11. [d] Work done B to C = $P_2 (V_C - V_B)$

$$= \mu R (T_C - T_B)$$

$$= 6 \times R \times (2200 - 800)$$

$$= 6R \times 1400$$

$$\text{Work done D to A} = P_1 (V_A - V_B)$$

$$= 6R \times (600 - 1200) = -6R (600)$$


Work done from A to B and B to C is zero because of a constant volume

$$\therefore \text{Total work done}$$

$$= 6R \times 1400 - 6R \times 600 = 6R (800)$$

$$= 50 \times 800 = 40000 \text{ J}$$

12. [a] Area under PV graph = work done



Here $\text{area}_2 > \text{area}_1 > \text{area}_3$

$$\therefore W_2 > W_1 > W_3$$

13. [d] Here $\frac{\pi p_1 A^4}{8\eta \ell} = \frac{\pi P_2 (2A)^4}{8\eta \ell}$

$$\text{i.e. } P_1 = 16P_2$$

$$\text{Given that } P_1 + P_2 = 1 \text{ m}$$

$$\therefore P_1 + \frac{P_1}{16} = 1$$

$$\text{or } P_1 = \frac{16}{17} = 0.94 \text{ m}$$

14. [b] Here $\left(\frac{T_1}{T_2}\right)^2 = \left(\frac{r_1}{r_2}\right)^3 = \left(\frac{r_1}{0.5r_1}\right)^3 = 8$

$$\text{or } \frac{T_1}{T_2} = 2\sqrt{2}$$

$$\text{i.e. } T_2 = \frac{T_1}{2\sqrt{2}} = \frac{365}{2\sqrt{2}} = 129 \text{ days}$$

15. [b] Power = $\frac{\text{work done}}{\text{time}}$; $P = \frac{vq}{t}$

$$\text{Here } V = mv = v \times 10^6$$

$$\text{and } q/t = Q$$

$$\therefore P = VQ \times 10^6$$

16. [a] $E_x = -\frac{dV}{dx} = \frac{4-2}{(6-4)10^{-2}} = -100 \text{ Vm}^{-1}$

$$E_y = \frac{dv}{dy} = \frac{2-4}{(2-1)10^{-2}} = 200 \text{ Vm}^{-1}$$

17. [b] Side of plate = \sqrt{A}

Capacitance of air portion

$$C_1 = \frac{\epsilon_0 \sqrt{A} (\sqrt{A} - x)}{d} = \frac{\epsilon_0 A - \epsilon_0 \sqrt{A} x}{d}$$

Capacitance = $C_1 + C_2$

$$= \frac{\epsilon_0 A}{d} - \frac{\epsilon_0 \sqrt{A} x}{d} + \frac{\epsilon_0 \epsilon_r \sqrt{A} x}{d}$$

$$= \frac{\epsilon_0}{d} (A - \sqrt{A} x + \epsilon_r \sqrt{A} x)$$

18. [d] Germanium is a semiconductor whereas copper is a metal conductor. A metal conductor has positive temperature coefficient of resistance whereas a semi-conductor has negative temperature coefficient of resistance. On cooling resistance of copper decreases whereas that of germanium increases.

19. [d] An ideal voltmeter has infinite resistance

$$\text{circuit current } I = \frac{60}{300 + 400} = \frac{3}{35} \text{ A}$$

$$\text{P.D. across } 400 \Omega \text{ is } \frac{3}{35} \times 400 = 34.3 \text{ V}$$

But the voltmeter reads 30 V across this resistor.

It simply mean that the voltmeter has error of 4.3 V


$$\text{PD across } 300 \Omega = 300 \times \frac{3}{35} = 25.7 \text{ V}$$

$$\text{Reading of same voltmeter} = (25.7 - 4.3) = 21.4 \text{ V}$$

20. [a] Here $I = \frac{2}{10+R}$

Potential difference across 40 cm wire = $I \times \text{resistance of 40 cm of wire}$

$$= \frac{2}{10+R} \times 40 \left(\frac{10}{100} \right) = \frac{8}{10+R}$$



But as per statement

$$\frac{8}{10+R} = 10 \text{ mV} = 10 \times 10^{-3} \text{ or } R = 790 \Omega$$

21. [a] Magnetic field due to ab is zero because O lies on the extended wire itself.

Magnetic field due to infinite wire cd is


$$B_1 = \frac{\mu_0}{4\pi r} (\sin 0^\circ + \sin 90^\circ) = \frac{\mu_0 i}{4\pi r}$$

Magnetic field due to circular portion

$$B_2 = \frac{\mu_0}{4\pi} \frac{i \left(\frac{3}{4} 2\pi r \right)}{r^2} = \frac{\mu_0 i}{4\pi r} \frac{3\pi}{2}$$

$$\therefore B = B_1 + B_2 = \frac{\mu_0 i}{4\pi r} \left(\frac{3}{2} \pi + 1 \right)$$

22. [d] Here $B_R = 2B \sin \alpha$ ($\because \cos$ component cancel out)



$$= \frac{\mu_0 I}{2\pi a} \left(\frac{d}{a} \right) = \frac{\mu_0 I d}{\pi r^2}$$

23. [b] Here $\tau = nIAB \sin \theta$, here θ is angle between magnetic induction and normal to the surface of loop.


$$\text{Here } A = \frac{1}{2} \times \text{base} \times \text{height}$$

$$= \frac{1}{2} \times 0.02 \times 0.02 \sin 60^\circ$$

$$= 1.732 \times 10^{-4}$$

$$= 8.66 \times 10^{-1} \text{ Nm}$$

24. [c] For a prism which in thin $\delta = (\mu - 1) A$



$$\text{Then } yy' = S \times \delta \left(\because \theta = \frac{\ell}{r} \right)$$

$$= S(\mu - 1)A = As(\mu - 1)$$

25. [a] Here $\theta = 60^\circ = \frac{\pi}{3}$

$$= \frac{1}{60} = \frac{\pi}{180} \times \frac{1}{60} \text{ rad.}$$

Distance between persons $\ell = 3 \text{ m}$ using

$$v = \frac{\ell}{r} \text{ we get } v = \frac{\ell}{x} = \frac{3}{x}$$

$$\text{or } x = \frac{3}{v} = \frac{3 \times 180 \times 60}{\pi} \text{ or } x = 10 \text{ km}$$

26. [d] Diffraction is observable if the width of slit is of the order of the wavelength of wave used. Since in the given problem wavelength of x-ray is too less than width of slit. So diffraction pattern will not be observed.

27. [b] Using $I = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \phi$ when

$$I_1 = I \text{ and } I_2 = 4I$$

$$\text{When } \phi = \frac{\pi}{2} \text{ (at A),}$$

$$I_A = I + 4I = 5I$$

$$\text{when } \phi = \pi \text{ (at B),}$$

$$I_B = I + 4I - 4I = I \therefore I_A - I_B = 4I$$

28. [a] Using $P = \sqrt{2mE}$, we get

$$\frac{P_\alpha}{P_p} = \sqrt{\frac{2 \times 4m_p \times E_\alpha}{2 \times m_p \times \epsilon_p}} = \sqrt{\frac{2 \times 4m_p \times E_\alpha}{2 \times m_p \times 16E_\alpha}} = \frac{1}{2}$$

$$\text{But } \frac{\lambda_p}{\lambda_\alpha} = \frac{\lambda_\alpha}{P_p} \therefore \frac{\lambda_p}{\lambda_\alpha} = \frac{1}{2}$$

29. [b] Using $E = \frac{nhc}{\lambda}$ we get

$$10^{-7} = \frac{n(6.6 \times 10^{-34})(3 \times 10^8)}{(5000 \times 10^{-10})}$$

$$n = 2.5 \times 10^{11}$$

30. [a] Using Moseley's law for $K\alpha$ line, we get

$$\frac{1}{\lambda} = \frac{3}{4} R(Z-1)^2$$

$$\text{or } \frac{1}{0.76 \times 10^{-10}} = \frac{3}{4} (1.09 \times 107) (z-1)^2$$

$$\text{or } \frac{4 \times 10^3}{0.76} = 3 \times (1.09) (z-1)^2$$

$$\text{or } (z-1)^2 = \frac{4 \times 10^3}{0.76 \times 3 \times 1.09} \approx 1000$$

$$\text{or } Z-1 = 40$$

$$\text{or } Z = 41$$

CHEMISTRY

31. [d] Meq. of Acid = Meq. of $\text{Ba}(\text{OH})_2$

$$\Rightarrow \frac{1.25}{M/2} \times 1000 = (0.25 \times 2) \times 25$$

$$\Rightarrow M = 200$$

32. [b] $\frac{r_{(H_2)}}{r_{(D_2)}} = \sqrt{\frac{M_{(D_2)}}{M_{(H_2)}}} = \sqrt{\frac{4}{2}} = \frac{\sqrt{2}}{1}$

33. [a] From $K_p = K_c (\text{RT})^{\Delta n_g}$
 $= 1.8 \times 10^{-4} \times (0.082 \times 298)^2 = 0.108$

34. [a] $\Delta H_{\text{sublimation}} = \Delta H_{\text{fusion}} + \Delta H_{\text{vap}}$

35. [d] $K_{sp} = 4s^3 = 4 \times 10^{-12} \Rightarrow s = 10^{-4} \text{ M}$

36. [a] Oxidant is the one whose O.N. decreases during the reaction. H_2SO_4 (O.N. of S = +6) changes to SO_2 (O.N. of S = +4)

37. [d] $d = \frac{Z \times M}{a^3 \times N_A} = \frac{2 \times (3 \times 10^{-3})}{\left(\frac{4 \times 4.52 \times 10^{-3}}{2\sqrt{3}}\right)^3 \times 6.02 \times 10^{23}}$
 $= 900 \text{ kg m}^{-3}$




38. [b] $E_{\text{cell}} = E_{\text{cell}}^0 + \frac{0.059}{n} \log \frac{[\text{Cathode}]}{[\text{Anode}]}$
 $= [-0.0403 - (-0.763)] + \frac{0.059}{2} \log \frac{0.004}{0.2}$
 $= +0.36 + \frac{0.059}{2} \log \frac{0.04}{2}$

39. [d] $\frac{\Delta P}{P^0} = \frac{W_B M_A}{M_B W_A}$ or $M_B = \frac{W_B M_A}{W_A (\Delta P / P^0)}$
 $\Rightarrow M_B = \frac{2.5 \times 78 \times 640}{39 \times 40} = 80$



40. [a] $t = \frac{2.303}{k} \log \frac{a}{a-x} = \frac{2.303}{6} \log \frac{0.5}{0.05}$
 $= 0.384 \text{ min}$

41. [d] Isocyanide test also known as carbylamine test.

42. [a] 4-methyl benzene sulphonic acid is stronger than acetic acid thus it will release acetic acid from sodium acetate.




46. [a] Libermann's reaction




49. [a] Free rotation around carbon-carbon bond takes place easily in alkanes. Now ethane and hexachloroethane both are alkanes, but in hexachloroethane bulky chlorine atom is present while ethane is least hindered.


50. [c] Due to the presence of $-\text{Cl}$ group which is a $+\text{M}$ group.

51. [a] Due to similar charges on adjacent atom the structure is least stable.






Free radical is formed which is sp^2 hybridised



53. [a] If atom or group of higher priority are on opposite direction at the double bond of each carbon atom then the configuration is known as E and if they are in same direction then the configuration is known as Z-configuration.



54. [c] The brown ring test for NO_2^- and NO_3^- is due to formation of $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+}$

55. [a] The absorption of energy or observation of color in a complex transition compounds depend upon the charge of the metal ion and the nature of the ligand attached. The same metal ion with different ligands shows different absorption depending upon the type of ligand, the presence of weak field ligand make the central metal ion to absorb low energies i.e. of higher wavelength.

56. [c] The existence of Fe^{2+} and NO^+ in nitroprusside ion $[\text{Fe}(\text{CN})_5\text{NO}]^{2-}$ can be established by measuring the magnetic moment of the solid compound which should correspond to $\text{Fe}^{2+} = 3d^6$ four unpaired electron.

57. [b] LiNO_3 on heating gives



58. [d] Three dimensional sheet structure are formed when three oxygen atoms of each $[\text{SiO}_4]^{4-}$ tetrahedral are shared.

59. [a] 'Lapis Lazuli' is the aluminium silicate present in the earth rocks as blue stone.

60. [b] $B < C < N < O$ when we move from B to O in a periodic table the first ionization enthalpy increase due to the attraction of nucleus towards the outer most of electron and IE of $N > O$.

MATHEMATICS

61. [d] Mean = $\frac{\sum x_i w_i}{\sum w_i} = \frac{0C_0 + 1C_1 + \dots + nC_n}{C_0 + C_1 + \dots + C_n}$
 $= \frac{n \cdot 2^{n-1}}{2^n}$

62. [a] $f(x) = 1 + x \sin x [\cos x]$

$$\because 0 < x \leq \frac{\pi}{2} \Rightarrow 0 \leq \cos x < 1$$

$$\Rightarrow [\cos x] = 0$$

$$\therefore f(x) = 1$$

$\therefore f(x)$ is a constant function and hence continuous. It neither strictly increasing nor decreasing.

63. [b] It is given that $\left(\frac{\delta r}{r}\right) \times 100 = 1$

$$V = \frac{4}{3} \pi r^3 \Rightarrow \log V = \log \frac{4\pi}{3} + 3 \log r$$

$$\frac{1}{V} \delta V = \frac{3}{r} \delta r$$

$$\frac{\delta V}{V} \times 100 = \frac{3\delta r}{r} \times 100$$

$$= 3 \times 1 = 3$$

Hence error in volume is with in 3%

64. [c] Set $A = \{1, 2, 3\}$

$$\text{and } R = \{(1, 1), (2, 2)\}$$

Since $(3, 3) \notin R$ it is not reflexive

$$\text{Since } R^{-1} = \{(1, 1), (2, 2)\} = R,$$

R is symmetric

Since the situation in $(a, b), (b, c) \in R$ does not arise in R , R is also transitive.

$$\text{Also } R \cap R^{-1} = \{(1, 1), (2, 2)\} \subset D_A$$

$$= \{(1, 1), (2, 2), (3, 3)\}$$

$\Rightarrow R$ is anti symmetric

Hence (ii) (iii) and (iv) are correct.

65. [c] $\{(A - B) \cup (B - C) \cup (C - A)\}^c$
 $= (A \cup B \cup C) - \{(A - B) \cup (B - C) \cup (C - A)\}$
 $= A \cap B \cap C \{ \because A \cup B \cup C = \text{universal set} \}$

66. [c] $(B^{-1}AB)^2 = (B^{-1}AB)(B^{-1}AB)$

$$= (B^{-1}ABB^{-1}AB)$$

$$= (B^{-1}A^2B)$$

$$= (B^{-1}A^2B)$$

$$(B^{-1}AB)^3 = (B^{-1}AB)^2 (B^{-1}AB)$$

$$= (B^{-1}A^2B)(B^{-1}AB)$$

$$= (B^{-1}A^2BB^{-1}AB)$$

$$= (B^{-1}A^2IAB)$$

$$= (B^{-1}A^3B)$$

$$\begin{aligned} \text{Now } (B^{-1}AB)^4 &= (B^{-1}AB)^3 (B^{-1}AB) \\ &= (B^{-1}A^3B) (B^{-1}AB) \\ &= B^{-1}A^4B \end{aligned}$$

67. [a] $\therefore g(0) = 0 \quad \therefore \lim_{x \rightarrow 0} \frac{g(x)}{x} \left(\frac{0}{0} \right)$ form
 $\therefore \lim_{x \rightarrow 0} g'(x) = g'(0) \quad \dots (1)$

$$\therefore g(x) = \begin{vmatrix} f(x+\alpha) & f(x+2\alpha) & f(x+3\alpha) \\ f(\alpha) & f(2\alpha) & f(3\alpha) \\ f'(\alpha) & f'(2\alpha) & f'(3\alpha) \end{vmatrix}$$

$$\therefore g'(x) = \begin{vmatrix} f'(x+\alpha) & f'(x+2\alpha) & f'(x+3\alpha) \\ f(\alpha) & f(2\alpha) & f(3\alpha) \\ f'(\alpha) & f'(2\alpha) & f'(3\alpha) \end{vmatrix}$$

$$\therefore g'(0) = 0$$

$$\lim_{x \rightarrow 0} \frac{g(x)}{x} = g'(0) = 0$$

68. [a] $\Delta_1 = \begin{vmatrix} f & 2d & e \\ 2z & 4x & 2y \\ e & 2a & b \end{vmatrix} = - \begin{vmatrix} 2d & f & e \\ 4x & 2z & 2y \\ 2a & e & b \end{vmatrix} (C_1 \leftrightarrow C_2)$

$$= \begin{vmatrix} 2d & e & f \\ 4x & 2y & 2z \\ 2a & b & e \end{vmatrix} (C_2 \leftrightarrow C_3)$$

$$= - \begin{vmatrix} 2a & b & e \\ 4x & 2y & 2z \\ 2d & e & f \end{vmatrix} (R_1 \leftrightarrow R_3)$$

$$= \begin{vmatrix} 2a & b & e \\ 2d & e & f \\ 4x & 2y & 2z \end{vmatrix} = \Delta_2 (R_2 \leftrightarrow R_3)$$

$$\therefore \Delta_1/\Delta_2 = 1$$

69. [d] \therefore All coins are identical

\therefore First we will give 3 coin to each person so that every one has at least 3 rupee, now rest 5 coin we have to distribute among 5 person in such a way that any one can get any no. of coin.

\therefore Total no. of ways

$${}^{5+5-1}C_{5-1} = {}^9C_4 = 126$$

$\{\therefore$ No. of ways of distributing n identical thing among r person when any one can get any no. of thing is ${}^{n+r-1}C_{r-1}$

70. [c] $x_1 < x_2 \leq x_3 < x_4 < x_5 \leq x_6$ gives rise to the following four cases

$$x_1 < x_2 < x_3 < x_4 < x_5 < x_6$$

$$x_1 < x_2 = x_3 < x_4 < x_5 < x_6$$

$$x_1 < x_2 < x_3 < x_4 < x_5 = x_6$$

$$x_1 < x_2 = x_3 < x_4 < x_5 = x_6$$

$$\therefore \text{Total ways } {}^9C_6 + {}^9C_5 + {}^9C_5 + {}^9C_4$$

$$= {}^{10}C_6 + {}^{10}C_5 = {}^{11}C_6$$

71. [d] $2 \left\{ 1 + \frac{a^2}{2} + \frac{a^4}{4} + \dots \right\} = e^a + e^{-a}$

$$= e^{ln n} + e^{-ln n} = n + \frac{1}{n} = \frac{n^2 + 1}{n}$$

72. [a] $\left(\sqrt{x} - \frac{2}{x} \right)^{18}$

Let (r+1)th term is independent of x

$$\therefore r = \frac{18 \times \frac{1}{2} - 0}{\frac{1}{2} + 1} = 6 \quad \{ \therefore r = \frac{n\alpha - m}{\alpha + \beta} \}$$

$\therefore (r+1) = 7^{\text{th}}$ term is independ of x

$$\therefore 7^{\text{th}} \text{ term is } {}^{18}C_6 (\sqrt{x})^{18-6} \left(-\frac{2}{x} \right)^6$$

$$= {}^{18}C_6 2^6$$

73. [c] $S = 1 + \frac{2}{3} \cdot \frac{1}{2} + \frac{2.5}{3.6} \left(\frac{1}{2} \right)^2 + \frac{2.5.8}{3.6.9} \left(\frac{1}{2} \right)^3 + \dots$

$$= 1 + \frac{2/3}{1} \left(\frac{1}{2} \right) + \frac{(2/3)(5/3)}{2} \left(\frac{1}{2} \right)^2$$

$$+ \frac{(2/3)(5/3)(8/3)}{3} \left(\frac{1}{2} \right)^3 + \dots$$

$$= \left(1 - \frac{1}{2} \right)^{-2/3} = \left(\frac{1}{2} \right)^{-2/3} = 2^{2/3} = 4^{1/3}$$

74. [d] $\therefore \omega = \frac{-1 + i\sqrt{3}}{2} \Rightarrow i\omega = \frac{-\sqrt{3} - i}{2}$

$$\therefore \arg(i\omega) = \pi + \frac{\pi}{6}$$

$$\therefore \omega^2 = \frac{-1 - i\sqrt{3}}{2} \Rightarrow i\omega^2 = \frac{\sqrt{3} - i}{2}$$

$$\therefore \arg(i\omega^2) = 2\pi - \frac{\pi}{6}$$

$$\therefore \arg i\omega + \arg i\omega^2 = 3\pi$$

75. [a] $\therefore \frac{1}{\log_2 4} = \frac{1}{\frac{1}{r} \log_2 4} = \frac{r}{2}$

$$\therefore \sum_{r=1}^n \frac{r}{2} = \frac{1}{2} \left(\frac{n(n+1)}{2} \right) = \frac{n(n+1)}{4}$$


76. [c] $a_1 + a_8 + a_{15} = 3a_1 + 21d = 15$
 $\Rightarrow a_1 + 7d = 5$
 $a_2 + a_3 + a_8 + a_{13} + a_{14}$
 $= 5a_1 + 35d$
 $= 5(a_1 + 7d)$
 $= 5 \times 5 = 25$

77. [c] $f(x) = (x - a_1)(x - a_3)(x - a_5) + 2(x - a_2)(x - a_4)(x - a_6) = 0$
 $a_1 < a_2 < a_3 < a_4 < a_5 < a_6$
 $f(a_1) = 2(a_1 - a_2)(a_1 - a_4)(a_1 - a_6) < 0$
 $f(a_2) = (a_2 - a_1)(a_2 - a_3)(a_2 - a_5) > 0$
 \therefore At least one real root lies in (a_1, a_2)
 Similarly, at least one real roots lies in each interval (a_3, a_4) and (a_5, a_6)
 But $f(x)$ is cubic, therefore there are only three roots.
 Hence the equation $f(x) = 0$ has one real roots in each interval (a_1, a_2) , (a_3, a_4) and (a_5, a_6) .

78. [a] $x dx + z dy + (y + 2z) dz = 0$
 $\Rightarrow x dx + 2z dz + z dy + y dz = 0$
 $x dx + 2z dz + d(yz) = 0$
 $\frac{x^2}{2} + z^2 + yz = c$

79. [a] Slope of tangent
 $\frac{dy}{dx} = 2x + 1$
 $\Rightarrow y = x^2 + x + C$
 when $x = 1, y = 2$
 $\therefore 2 = 1 + 1 + C \Rightarrow C = 0$
 $\therefore y = x^2 + x$
 \therefore Required area $= \int_0^1 (x^2 + x) dx$
 $= \left[\frac{x^3}{3} + \frac{x^2}{2} \right]_0^1 = \frac{5}{6}$

80. [b]




By symmetry
 Let $P = (t, 0)$ then
 $Q(t, e^{-t}), R = (-t, e^{-t})$ and $S = (-t, 0)$
 \therefore Area of rectangle $= 2te^{-t} = f(t)$ say

then $\frac{df}{dt} = 2\{-te^{-t} + e^{-t}\} = 0 \Rightarrow t = 1$
 $\frac{d^2f}{dt^2} = 2\{-(1-t)e^{-t} - e^{-t}\} < 0$ for $t = 1$
 Hence, maximum area $2/e$

81. [c] $I = \int |\sin x| dx$
 $= \begin{cases} \int \sin x dx & \text{if } \sin x \geq 0 \\ -\int \sin x dx & \text{if } \sin x < 0 \end{cases}$
 $= -\cos x + C$ if $\sin x \geq 0$
 $\cos x + C$ if $\sin x < 0$
 $= \cos x \cdot \text{sgn}(\sin x) + C$
 $\{\because \text{sgn}(\sin x) = \frac{|\sin x|}{\sin x}$
 $= \begin{cases} 1 & ; \sin x > 0 \\ 0 & ; \sin x = 0 \\ -1 & ; \sin x < 0 \end{cases}$

82. [b] Feasible region of constraints is shown in graph



Extreme points are, $(0, 0)$, $(15, 0)$, $(0, 12)$ and $(12, 6)$.

$\therefore P = 6x + 8y$
 $\therefore P_{(12, 6)} = 72 + 48 = 120$
 $P_{(0, 0)} = 0, P_{(15, 0)} = 90, P_{(0, 12)} = 16$
 Clearly maximum P is 120 at $(12, 6)$.
 Hence (b) is the correct answer.

83. [d] $\frac{dy}{dx} = -\frac{c^2}{x^2}$
 $\Rightarrow \left(\frac{dy}{dx}\right)_{(x', y')} = \frac{-c^2}{(x')^2} = -\frac{x' y'}{(x')^2} = -\frac{y'}{x'}$
 \therefore Equation of tangent at (x', y') is
 $y - y' = -\frac{y'}{x'}(x - x')$

Which meets the co-ordinate axes at A and B (say) then $A = (2x', 0)$, $B = (0, 2y')$
Mid point of AB is (x', y')


84. [d] D.R.'s of OP = a, a, a
 \therefore Equation of plane \perp to OP and passing through P is
 $a(x - a) + a(y - a) + a(z - a) = 0$
 $\Rightarrow x + y + z = 3a$
 Intercepts on axes made by the planes are
 $3a, 3a, 3a$
 \therefore Sum of reciprocal of the intercepts
 $= \frac{1}{3a} + \frac{1}{3a} + \frac{1}{3a} = \frac{1}{a}$

85. [c] $|\vec{a}|^2 = p^2 + 25 + 289 = p^2 + 314$
 $|\vec{b}|^2 = 4q + 169 + 1 = 4q + 170$
 According to question
 $|\vec{a}|^2 = |\vec{b}|^2$
 $\Rightarrow p^2 + 314 = 4q + 170$
 $\Rightarrow p^2 = 4q - 144 = 4(q - 36)$
 p, q are +ve integer
 $1 \leq p, q \leq 1000$
 p is even integer let $p = 2K$
 then $4K^2 = 4(q - 36)$
 $\Rightarrow K^2 = q - 36$
 $\therefore 1 \leq K^2 \leq 964$
 $\Rightarrow 1 \leq K \leq 31$
 \therefore Number of ordered pairs (p, q) = 31

86. [d] $\frac{x^2}{8-a} + \frac{y^2}{a-2} = 1$ will represent an ellipse
 if $8 - a > 0$, $a - 2 > 0$
 and $8 - a \neq a - 2$
 $\Rightarrow a < 8$, $a > 2$ and $a \neq 5$
 $\therefore a \in (2, 8) - \{5\}$

87. [d] $y = x - 1$ is a focal chord of the parabola $y^2 = 4x$. Therefore tangent at its extremities are perpendiculars.


88. [d]



$A_1B_1 = 2\sqrt{2}$
 $AB = 2\sqrt{2} - 2$
 $= 2(\sqrt{2} - 1)$
 $OA = \sqrt{2} - 1$
 taking origin as centre and OA as radius circle will touch all four circles

\therefore equation of circle is
 $x^2 + y^2 = (\sqrt{2} - 1)^2$
 $x^2 + y^2 = 3 - 2\sqrt{2}$

89. [a]



Clearly $a > 0$
 Also P lies on that side of line $x + y = 2$ where origin lies
 $\therefore a + a^2 - 2 < 0$
 $\Rightarrow (a - 1)(a + 2) < 0$
 $\Rightarrow -2 < a < 1$
 but $a > 0$
 $\therefore 0 < a < 1$
 $\therefore a \in (0, 1)$

90. [b] Let $a = 3K$, $b = 7K$ and $c = 8K$

$$\therefore s = \frac{a + b + c}{2} = 9K$$


$$\text{there } \frac{R}{r} = \frac{abc}{4\Delta} \cdot \frac{s}{\Delta} = \frac{abc s}{4s(s-a)(s-b)(s-c)}$$

$$= \frac{3K \cdot 7K \cdot 8K}{4 \cdot 6K \cdot 2K \cdot K} = \frac{7}{2} \Rightarrow \frac{R}{r} = \frac{7}{2}$$

91. [b] $y = f(x) = \frac{x}{1 + x \tan x}$

$$\Rightarrow \frac{dy}{dx} = \frac{1 + x \tan x - x(\tan x + x \sec^2 x)}{(1 + x \tan x)^2}$$

$$= \frac{1 - x^2 \sec^2 x}{(1 + x \tan x)^2}$$



$$\frac{dy}{dx} = 0 \Rightarrow x^2 = \cos^2 x$$

There is only one point in $\left(0, \frac{\pi}{2}\right)$ say x_1 , at

$$\text{which } \frac{dy}{dx} = 0$$

$$\text{at } x_1 - h \Rightarrow x^2 < \cos^2 x \quad \therefore \frac{dy}{dx} > 0$$

$$\& \text{ at } x_1 + h \Rightarrow x^2 > \cos^2 x \quad \therefore \frac{dy}{dx} < 0$$

\therefore at x_1 slope change from +ve to -ve

\therefore There is only one critical point in $\left(0, \frac{\pi}{2}\right)$ at which $f(x)$ has local maxima.

92. [b] $3\cos^2\theta - 2\sqrt{3}\sin\theta\cos\theta + 3\sin^2\theta = 0$
 $\Rightarrow (\sqrt{3}\cos\theta + \sin\theta)(\cos\theta - \sqrt{3}\sin\theta) = 0$
 $\Rightarrow \tan\theta = \frac{1}{\sqrt{3}}$ or $\tan\theta = -\sqrt{3}$
 $\therefore \theta = n\pi + \frac{\pi}{6}$ or $\theta = n\pi - \frac{\pi}{3}$
 $\therefore |r - s| = |-3 - 6| = 9$

93. [b] $\cot^{-1}\frac{n}{\pi} > \frac{\pi}{6}$
 $\Rightarrow \frac{\pi}{6} < \cot^{-1}\frac{n}{\pi} < \pi, n \in \mathbb{N}$
 $\{\because \cot^{-1}x \in (0, \pi)\}$
 $\Rightarrow -\infty < \frac{n}{\pi} < \sqrt{3}$
 $-\infty < n < \sqrt{3}\pi$
 $-\infty < n < 5.4$
 $\Rightarrow \max. n = 5 \quad \{\because n \in \mathbb{N}\}$

94. [a] $\therefore \tan \pi [x] = 0 \forall x \in \mathbb{R}$ since $[x] \in \mathbb{Z}$
 Period of $\{x\} = 1$
 \Rightarrow Period of $\sin 3\pi\{x\} = 1$
 Hence period of $f(x) = 1$

95. [c] $f(x) = \cos^{-1}\sqrt{\log_{[x]}\left(\frac{|x|}{x}\right)}$
 For domain $\frac{|x|}{x} > 0$
 $\Rightarrow x \in (0, \infty)$
 and $[x] > 0$ and $[x] \neq 1$
 $\Rightarrow x \geq 2 \quad \therefore x \in [2, \infty)$
 $\Rightarrow \frac{|x|}{x} = 1$ then $\log_{[x]}\left(\frac{|x|}{x}\right) = 0$
 $f(x) = \cos^{-1}0 = \frac{\pi}{2}$

96. [c] $\therefore f(a) = 0$
 $\therefore \lim_{x \rightarrow a} \frac{\log_e\{1 + 6f(x)\}}{3f(x)} \left(\frac{0}{0}\right)$ form
 $\Rightarrow \lim_{x \rightarrow a} 2 \times \frac{\log_e\{1 + 6f(x)\}}{6f(x)} = 2 \times 1 = 2$
 $\therefore \lim_{x \rightarrow 0} \frac{\log_e\{1 + x\}}{x} = 1$

97. [d] $\lim_{x \rightarrow 5^+} \frac{x^2 - 9x + 20}{x - [x]}$
 $= \lim_{h \rightarrow 0} \frac{(5+h)^2 - 9(5+h) + 20}{5+h - [5+h]}$
 $= \lim_{h \rightarrow 0} \frac{h^2 + h}{h} = 0$
 $\lim_{x \rightarrow 4^-} \frac{x^2 - 9x + 20}{x - [x]}$
 $= \lim_{h \rightarrow 0} \frac{(4-h)^2 - 9(4-h) + 20}{4-h - [4-h]}$
 $= \lim_{h \rightarrow 0} \frac{h^2 + h}{1-h} = 0$
 $\therefore P = 0$

98. [a] $(a + bx)e^{y/x} = x \quad \dots (1)$
 Differentiating, w.r.t. x we get
 $be^{y/x} + (a + bx)e^{y/x} \cdot \left(\frac{xy_1 - y}{x^2}\right) = 1$
 $\Rightarrow be^{y/x} + x \cdot \left(\frac{xy_1 - y}{x^2}\right) = 1 \quad \{\because (a + bx)e^{y/x} = x\}$
 $\Rightarrow bxe^{y/x} + xy_1 - y = x$
 $\Rightarrow xy_1 - y = x - bxe^{y/x}$
 $\Rightarrow xy_1 - y = ae^{y/x} \quad \dots (2)$ (from (1))
 $\Rightarrow xy_2 + y_1 - y_1 = ae^{y/x} \left[\frac{xy_1 - y}{x^2}\right]$
 $\Rightarrow x^3y_2 = ae^{y/x}(xy_1 - y) = (xy_1 - y)^2$ (from (2))
 $\Rightarrow \frac{1}{y_2}(xy_1 - y)^2 = x^3$

99. [b] $\int_{-2}^3 f(x)dx = \int_{-2}^{-1} f(x)dx + \int_{-1}^0 f(x)dx + \int_0^1 f(x)dx$
 $+ \int_1^2 f(x)dx + \int_2^3 f(x)dx$
 $= (-2)^3 + (-1)^3 + 0^3 + 1^3 + 2^3 = 0$

100. [b] $x^2f(x) + f\left(\frac{1}{x}\right) = 2$
 $I = \int_{1/3}^3 f(x)dx$ put $x = \frac{1}{t}, dx = -\frac{1}{t^2}dt$
 $\Rightarrow I = -\int_3^{1/3} f\left(\frac{1}{t}\right) \cdot \frac{1}{t^2} dt = \int_{1/3}^3 f\left(\frac{1}{x}\right) \cdot \frac{1}{x^2} dx$
 $\Rightarrow 2I = \int_{1/3}^3 \left(f(x) + \frac{1}{x^2}f\left(\frac{1}{x}\right)\right) dx$

$$= \int_{1/3}^3 \left[x^2 f(x) + f\left(\frac{1}{x}\right) \right] \frac{1}{x^2} dx = \int_{1/3}^3 \frac{2}{x^2} dx$$

$$= -2 \left[\frac{1}{x} \right]_{1/3}^3 = -2 \left[\frac{1}{3} - 3 \right] = \frac{16}{3} \Rightarrow I = \frac{8}{3}$$

ENGLISH PROFICIENCY

101. [b] Geraff :

Incorrect spelling.

- 'e' should be replaced with 'i'
- The word should end with 'e' after 'ff'

Giraffe :

Correct spelling.

Giraf :

'fe' is to be added in the end.

Gerraffe :

- 'Ge' is to be replaced with 'Gi' to make the correct spelling.

102. [a] Luxurious : (Plush)

Something full of all 'amenities' making life 'cozy' and 'snug'.

Delicious : Irrelevant as it means 'something very tasty.'

Comforting : 'Irrelevant' as it means 'giving necessary comforts', whereas 'Plush' means more than comforts.

Tasty : (Irrelevant)

It means 'delicious'

103. [a] Lively : Correct synonym to 'sprightly' as both means, 'someone dashing / energetic / enthusiastic'.

Beautiful : (Irrelevant)

Sportive : (Irrelevant)

Intelligent : (Irrelevant)

104. [d] Deadly : It means 'Fatal'.

Hence, this is not a proper antonym to 'innocuous'.

Ferocious : It means 'horrible'

Hence, irrelevant to the opposite of 'innocuous'.

Poisonous : It means 'venomous'.

Hence, an irrelevant 'antonym'.

Harmful : It is a perfect antonym of innocuous which itself means 'harmless'.

105. [d] Corruption :

Irrelevant

Worldliness :

Irrelevant

Favouritism :

Irrelevant

Nepotism : (Correct Answer) because

It's a kind of corruption in which the authority in power takes the advantage of giving opportunity to their relatives in their self interest.

106. [b] Cross : (to pass by, to intersect)

It means different

Hence, irrelevant.

Shuttle : (Proper answer)

It's a kind of "regular beats" of an air flight or bus service between the two stations.

Travel : It means to journey.

Hence, irrelevant.

Run : (to move regularly)

Hence, irrelevant.

107. [c] Sentence 1 is correct :

This option is wrong because the word 'ingenuous' means 'frank and simple' which is inappropriate.

Sentence 2 is correct :

This option is also wrong because the word 'ingenious' means 'clever or prudent' and this is inappropriate.

Both the words, i.e. 'ingenuous' and 'ingenious' if interchanged together respectively, it really makes both the sentences meaningful.

Hence, appropriate option.

Both the sentences can't be interchanged.

This is an incorrect option because words have been misinterpreted together.

Incorrect option.

108. [c] Far off :

It can't be used in place of 'aloof' as far off means long-long ago.

Hence, incorrect alternative .

Introvert : It means 'self-centred',

Hence, It is an incorrect alternative.

distance : This is an appropriate word because one of the meaning of 'aloof' is distant also while keeping distance between two nouns.

Depressed : (it means 'hopeless')

Hence, quite irrelevant.

109. [a] "Meatless days" This is the name of a novel. Hence, no error is there.

Have been made : (Erroneous)

Because 'have' should be replaced with 'has' because 'meatless days' is a singular noun.

Into a film :

No error in this part of the sentence.

No error : Incorrect option because there is an error in the sentence.

110. [a] **Broke out :** (to start suddenly)

'Correct and relevant' option because it is used for 'wars' and 'diseases' e.g. cholera broke out in Surat in 1985.

Set out : (to start)

it is different because it is used when one leaves for somewhere

e.g. He set out on his long voyage to Achilrese.

took out : (incorrect use)

Because it means differently.

e.g. He took out a one rupee coin to give to the beggar.

Went out : (Incorrect use) Because meaning is different

e.g. : The light went out when I was preparing for my Board Exams.

Hence, inappropriate option.

LOGICAL REASONING

111. [d] The pattern is $x^2 + 1, x^2 + 2, \dots$

Missing number = $28 \times 2 + 3 = 59$

112. [a] A car runs on petrol and a television works by electricity.


113. [a] All except Titans are planets of the solar system.

116. [d]



117. [b] The third figure in each row comprises of parts which are not common to the first two figure.


119. [c]



120. [a]



121. [b]




122. [b] By taking reverse of the given letter and then by adding 1 in their number place.

$$\begin{aligned} ZIP &= (Z + I + P) \times 6 \\ &= (2 + 19 + 12) \times 6 = 198 \\ VIP &= (V + I + P) \times 6 \\ &= (6 + 19 + 12) \times 6 = 222 \end{aligned}$$

123. [a] One who travels in a train is called passenger
One who travels is Aeroplane is called Aeronaut

124. [c] All except Stone are obtained directly or indirectly from trees.

125. [b]




126. [b] Total letters = 12

group \Rightarrow 3 each

abb/bba/abb/bba

127. [a]



(a) (b) (c) (d)


✓

⊕ will be opposite of =

■ will be opposite of □

● will be opposite of ×

128. [c] Every thing is composed of molecules. Sun is different from Moon.



129. [b] $(1 \times 2 \times 5 \times 3) + (1 + 2 + 5 + 3) = 41$

$(4 \times 3 \times 6 \times 2) + (4 + 3 + 6 + 2) = 159$

$(5 \times 7 \times 3 \times 2) + (5 + 7 + 3 + 2) = 227$

130. [b]

